



Live Well, Work Well

Health and wellness tips for your work, home and life—brought to you by the insurance specialists at VistaNational Insurance Group, Inc.

While ASDs affect all groups of people, it has been found that they are nearly five times more common in boys than girls

CHILDREN'S HEALTH: AUTISM SPECTRUM DISORDERS

Autism spectrum disorders (ASDs) refer to a group of developmental disabilities that vary in severity and affect socialization, communication and other behaviors. The three types of ASDs are autistic disorder, Asperger syndrome and Pervasive Developmental Disorder-Not Otherwise Specified (PDD-NOS). Recent studies show that roughly 1 in 110 children have an ASD.

What are the ASDs?

Autistic disorder is characterized by impaired social interaction, communication issues, including problems with language, and unusual, repetitive or severely limited activities and interests. People with this disorder also often have intellectual disabilities.

Individuals with Asperger syndrome often experience some of the symptoms of autistic disorder, but less severe. While problems socializing and unusual behaviors and interests are not uncommon, Asperger syndrome is not associated with language difficulties or intellectual disability.

PDD-NOS is a term used to describe individuals who display only some of the signs of autistic disorder or Asperger syndrome. These individuals' symptoms are often milder and only affect socialization and communication.

ASDs are not mental illnesses or the result of bad parenting, and children with these disorders are not simply unruly kids who choose to misbehave.

Causes

While researchers are searching for answers, the exact causes of ASDs remain unknown.

Genetics

In many families, there appear to be patterns of ASDs,

supporting a genetic basis to the disorder. While no one gene has been identified as causing ASDs, researchers are searching for irregular segments of genetic code that children with these disorders may have inherited.

It also appears that some children are born with a susceptibility to ASDs, but a single "trigger" that causes the disorders to develop has yet to be identified. Other research efforts are investigating the possibility that under certain conditions, a cluster of unstable genes may interfere with brain development.

Environmental Factors

Another possible cause of ASDs is problems during pregnancy or delivery, particularly environmental factors like viral infections, metabolic imbalances and exposure to environmental chemicals. Some harmful substances, when ingested during pregnancy, have been associated with an increased risk of ASDs.

The Vaccination Debate

The question of a relationship between vaccines and ASDs continues to be debated. According to the Centers for Disease Prevention (CDC), research so far has not supported a link between the two.

Signs & Symptoms

All children develop differently, but signs of delayed development are generally present by 18 months of age. If you suspect that your child has an ASD, discuss your concerns with your child's pediatrician, who can recommend further evaluation. An ASD may be present if your child:

- Doesn't babble or respond to his or her name by 12 months of age
- Doesn't gesture, such as point or wave, by 14 months of age
- Repeats words or phrases over and over
- Doesn't play "pretend" by 18 months of age
- Appears obsessive or easily upset by change
- Loses previously acquired language or social skills at any age, or is delayed in developing them



CHILDREN'S HEALTH: AUTISM SPECTRUM DISORDERS

Screening & Diagnosis

Your child's doctor should check for signs of developmental delays during regular well-child visits. If your child shows some signs of an ASD, you may be referred to a specialist who can perform a comprehensive diagnostic evaluation. Because the disorders vary widely in severity, making a diagnosis may be difficult. There aren't medical tests to pinpoint these disorders. A formal evaluation consists of the specialist observing your child and talking to you about how your child's social skills, language skills and behavior have developed and changed over time. To help reach a diagnosis, your child may undergo a number of developmental tests covering speech, language and psychological issues.

Although the signs of ASDs often appear by 18 months, the diagnosis sometimes isn't made until around age 2, when delays in language and social development may be more obvious. Early diagnosis is important because early intervention is associated with the best chance for significant improvement.

Treatment & Prognosis

There are currently no cures for ASDs. ASDs cannot be outgrown, but individuals can learn to function within the confines of their disorders, especially if treatment begins early. Preschool children who receive intensive, individualized behavioral interventions often show significant progress. Your doctor can help identify the best resources and treatment options for your child, including:

- Behavioral and communication therapies—Programs have been developed to address the range of social, language and behavioral difficulties associated with ASDs. Some programs focus on reducing problem behaviors and teaching new skills, while others focus on teaching children how to act in social situations or how to communicate better with others.
- Drug therapies—Right now, there are no medications that directly improve the core signs of ASDs, but some do help control obsessive and aggressive behaviors. However, certain medications are only considered after behavior management has failed.
- Complementary or alternative approaches—Supplement educational and medical intervention with complementary therapies, such as art therapy, music therapy, special diets, vitamin and mineral supplements, and sensory integration.

Children with ASDs often respond well to structured educational programs. Successful programs often include a team of specialists and a variety of activities to improve social skills, communication and behavior.